
MŰHELYTANULMÁNYOK

DISCUSSION PAPERS

MT-DP – 2012/29

Cooperation with Externalities and Uncertainty

HELGA HABIS - DÁVID CSERCSIK

**Discussion papers
MT-DP – 2012/29**

**Institute of Economics, Research Centre for Economic and Regional Studies,
Hungarian Academy of Sciences**

**KTI/IE Discussion Papers are circulated to promote discussion and provoke comments.
Any references to discussion papers should clearly state that the paper is preliminary.
Materials published in this series may subject to further publication.**

Cooperation with Externalities and Uncertainty

Author:

**Helga Habis
junior research fellow
Institute of Economics
Research Centre for Economic and Regional Studies
Hungarian Academy of Sciences
Email: habis.helga@krtk.mta.hu**

**Dávid Csercsik
postdoctoral assistant professor
Pázmány Péter Catholic University
Email: csercsik@scl.sztaki.hu**

October 2012

**ISBN 978-615-5243-32-5
ISSN 1785 377X**

Cooperation with Externalities and Uncertainty

Helga Habis - Dávid Csercsik

Abstract

We introduce a new class of cooperative games where the worth of a coalition depends on the behavior of other players and on the state of nature as well. We allow for coalitions to form both before and after the resolution of uncertainty, hence agreements must be stable against both types of deviations. The appropriate extension of the classical core concept, the Sustainable Core, is defined for this new setup to test the stability of allocations in such a complex environment.

A prominent application, a game of consumers and generators on an electrical energy transmission network is examined in details, where the power in- and outlets of the nodes have to be determined in a way, that if any line instantaneously fails, none of the remaining lines may be overloaded. We show that fulfilling this safety requirement in a mutually acceptable way can be achieved by choosing an element in the Sustainable Core.

Keywords: partition function form games, uncertainty, core, sustainability

JEL classification: C71, C73, D62, L14, L94

Kooperáció externáliák és bizonytalanság mellett

Habis Helga – Csercsik Dávid

Összefoglaló

Bevezetjük a kooperatív játékelmélet egy olyan új játékosztályát, amelyben egy koalíció értéke függ a többi játékos viselkedésétől és a világhállapottól is. Egy koalíció megalakulhat a bizonytalanság feloldása előtt és után is, így a megoldásnak mindkét típusú blokkolással szemben stabilnak kell lennie. Kiterjesztjük a klasszikus mag definícióját erre a komplex környezetre, definiáljuk a fenntartható magot (sustainable core).

Részletesen elemzünk egy fontos alkalmazást, amely az elektromos átviteli hálózatokat érinti. Ezekben az energiabetáplálást és -fogyasztást úgy kell meghatározni, hogy a rendszer még egy vezeték hibája esetén is stabil maradjon, azaz a megmaradt vezetékek nem lehetnek túlterheltek. Megmutatjuk, hogy ezen rendszerbiztonsági feltétel biztosítható, ha egy fenntartható magbeli elemet választunk. Ez a megoldás ráadásul a fogyasztók és a termelők számára is elfogadható elosztást biztosít.

Tárgyszavak: partíciós függvény alakú játékok, bizonytalanság, mag, fenntarthatóság

JEL kódok: C71, C73, D62, L14, L94

Cooperation with Externalities and Uncertainty

Helga Habis* and Dávid Csércsik †

November 5, 2012

Abstract

We introduce a new class of cooperative games where the worth of a coalition depends on the behavior of other players and on the state of nature as well. We allow for coalitions to form both before and after the resolution of uncertainty, hence agreements must be stable against both types of deviations. The appropriate extension of the classical core concept, the *Sustainable Core*, is defined for this new setup to test the stability of allocations in such a complex environment.

A prominent application, a game of consumers and generators on an electrical energy transmission network is examined in details, where the power in- and outlets of the nodes have to be determined in a way, that if any line instantaneously fails, none of the remaining lines may be overloaded. We show that fulfilling this safety requirement in a mutually acceptable way can be achieved by choosing an element in the Sustainable Core.

Keywords: *partition function form games, uncertainty, core, sustainability*

JEL Classification: *C71, C73, D62, L14, L94*

*Department of Economics, Lund University, and Department of Microeconomics, Corvinus University of Budapest, and Institute of Economics, Research Centre for Economic and Regional Studies, Hungarian Academy of Sciences. E-mail: helga.habis@uni-corvinus.hu.

†Pázmány Péter Catholic University. Email: csercsik@scl.sztaki.hu. Financial support from the Hungarian Scientific Research Fund (OTKA PD-101106 and OTKA NF-104706), from the Momentum Programme (LD-004/2010) of Hungarian Academy of Sciences and the Excellence in Research Award of Corvinus University of Budapest are gratefully acknowledged.

1 Introduction

Cooperative game theory analyzes how the gains of cooperation is to be divided among the members of a coalition. The most famous solution to this problem is the *core* (Gillies, 1959). In a classical, static cooperative game it is implicitly assumed that players can make binding agreements regarding the allocation of the value of a coalition if they cooperated. In most real life situations the core is not applicable, since its assumptions are too restrictive. At the same time cooperative game theory is often criticized for not being able to incorporate externalities appropriately. In this paper we introduce a new class of transferable utility games, where we allow for dynamics, uncertainty and externalities as well, and propose an extended notion of the core as a solution to this game which relaxes its restrictive assumptions.

Dynamic cooperation has already been studied by a number of papers. The introduction of dynamics only, already raises a crucial question; namely what are the consequences of a current deviation for the future, or more precisely, how to define a profitable deviation? Different answers to this question lead to different solution concepts; the Social Nash Optimum (Grossman, 1977), the core (Bester, 1984), the Segregated Core (Repullo, 1988), the Two-stage Core (Koutsougeras, 1998), the Strong Sequential Core (Predtetchinski, Herings, and Peters, 2002) and the Weak Sequential Core (Kranich, Perea, and Peters, 2005; Habis and Herings, 2010). Habis and Herings (2011a) give a comparison of all these concepts in a two-period general equilibrium setting and conclude that the Weak Sequential Core is the most satisfactory concept so far. Habis and Herings (2011b) introduce *transferable utility games with uncertainty*, define the Weak Sequential Core in this setup and give its characterization.

In many cases the value of a coalition may well depend on the behavior of the rest of the players, thus we may have externality among coalitions. This situations can be modeled by a *partition function form game*, introduced by Thrall and Lucas (1963). They take the pessimistic approach that residual players aim to minimize the payoff of the deviators. This idea is in the spirit of the α -core (Aumann and Peleg, 1960), defined for NTU-games, where it is assumed that a coalition deviates only if it gets a higher payoff irrespective of the induced partition of the rest of the players. Its complete opposite is the optimistic approach of the ω -core (Shenoy, 1980), where deviators expect to be helped by residuals. A more rational reaction is applied in the γ -core (Chander and Tulkens,

1997), where the deviating coalition must face individually best responses. Here it is true again, that different assumptions regarding residual behavior may lead to further different solution concepts; e.g. Hart and Kurz (1984); van der Laan and Cornet (1998); Huang and Sjöström (2003). Here we will concentrate on the Recursive Core by Kóczy (2007), since it is less sensitive to optimism or pessimism than the α - and ω -cores, furthermore allows a rational residual response as the γ -core, but this response is general, endogenous and consistent with the solution of the main game.¹

In this paper we will combine the above two directions of research, and allow for the presence of uncertainty and externalities at the same time by introducing the *partition function form game with uncertainty* (hereafter PFU-game). A PFU-game consists of two time periods, 0 and 1. In period 1 one out of a finite number of states of nature may materialize and conditional on the state, the players are involved in a particular partition function form game. An outcome therefore specifies a payoff-partition configuration conditional on each possible state of nature. A utility function is then used to assign a utility level to each profile of state-contingent payoffs.

We are interested in the appropriate definition of the core of a PFU-game. In this setting coalitions (forming some partition) are allowed to form in both time periods. Stability requires that a suggested allocation cannot be blocked by any coalition at any period, i.e. both before and after the resolution of uncertainty. Binding agreements are clearly not natural in such an environment, thus only self-enforcing agreements should be allowed for. These considerations lead to the concept of the *Sustainable Core*. Extending the characterization of the Weak Sequential Core (Habis and Herings, 2011b), we say that an allocation belongs to the Sustainable Core only if conditional on the state of nature it belongs to the Recursive Core of the PFF-game related to that state, and moreover there is no coalition (forming some partition) in period 0 that can propose state-contingent Recursive Core elements of the game restricted to that coalition, which gives its members higher utility.

An important application of the PFU-game is the electrical energy transmission network, where consumers and generators need to find a stable allocation of power in- and outlets, taking into account possible line failures. Cserecsik and Kóczy (2011) show that both negative and positive externalities exist in a static electrical energy transmission network game of generators and consumers, and therefore suggest the Recursive Core as a

¹Further desirable properties are discussed in (Kóczy, 2006).

solution concept. Here we extend their example by the possibility of line failures in the grid, which creates the uncertainty. We show that the Sustainable Core can be used to find a solution to this extended game.

The organization of the paper is as follows. First we introduce the notation and the definition of the core for the PFF-game in Section 2 followed by the definition of the PFU-game and our solution concept in Section 3. The application to the electrical energy transmission network is analyzed in Section 4 in detail, finally Section 5 concludes.

2 Preliminaries

Let $N = \{1, 2, \dots, n\}$ be the set of players, and its non-empty subsets are the coalitions, denoted by C . A cooperative game with transferable utility, or characteristic function form game, is a pair (N, v) , where $v : 2^N \rightarrow \mathbb{R}$ is a characteristic function which assigns to each coalition $C \subseteq N$ its *worth* $v(C)$, with the convention that $v(\emptyset) = 0$.

2.1 Partition Function Form Games

A *partition* P is a set of disjoint coalitions; $P = \{C^1, C^2, \dots, C^m\}$, where their union is N ; i.e., players in set K cooperate if and only if $K \in P$. The set of partitions is \mathcal{P} and the set of partitions of $C \in N$ is $\mathcal{P}(C)$, with a typical element P^C . A partition function; $w : \mathcal{P} \rightarrow (2^N \rightarrow \mathbb{R})$ assigns a characteristic function to each partition. A *cooperative game with transferable utility in partition function form*, or briefly *PFF-game*, is a pair (N, w) , and it is an extension of the characterization function form game where we allow for externalities across players and coalitions; hence, the worth of a coalition may be different in each partition. For $C \in P$, let the worth $w(C, P)$ denote the amount that the players in C can guarantee themselves by cooperating, when the coalition C is embedded in the partition P . An *outcome* is a pair $\omega = (x, P)$, consisting of a payoff vector $x = (x^1, \dots, x^n) \in \mathbb{R}^N$ satisfying feasibility; $\sum_{i \in (C \in P)} x^i \leq w(C, P)$ for all $C \in P$, and a partition $P \in \mathcal{P}$. The payoff for a coalition C is a vector $x^C = (x^i)_{i \in C} \in \mathbb{R}^C$. Let us denote the the set of outcomes in (N, w) by $\Omega(N, w)$.

In a PFF-game whether a coalition has a profitable deviation, depends on the induced partition of the players. There are a number of ways to model the reaction of the remaining

players, which lead to different solution concepts. Here we concentrate on the *Recursive Core* by Kóczy (2007), that allows the remaining, residual players to freely react, and form a core-stable partition before the payoff of the deviating coalition is evaluated. Thus, given a deviation, the residual players face the problem of solving another, smaller PFF-game. We call this a *residual game*.

Definition 2.1. Consider a game (N, w) and a player set $R \subsetneq N$. Assume $\overline{R} = N \setminus R$ have formed the partition $P^{\overline{R}} \in \mathcal{P}(\overline{R})$. Then the residual game $(R, w^{P^{\overline{R}}})$ is the PFF game over the player set R with the partition function given by $w^{P^{\overline{R}}}(C, P^R) = w(C, P^R \cup P^{\overline{R}})$.

Observe that the subset of players in the residual game still experience the externalities from the players in \overline{R} . Hence, if the core is the solution for (N, w) , then the core solves the residual game $(R, w^{P^{\overline{R}}})$ as well. The definition of the residual game is similar to that of the reduced game (Moulin, 1985), where the payoff structure of the larger game is accounted for, but the significance of the partition of \overline{R} is not considered.

Next we define the Recursive Core in the spirit of (Kóczy, 2007), by induction on the number of players.

Definition 2.2 (Recursive Core). The definition consists of three steps. Consider a PFF-game (N, w) .

1. *Trivial game* For a single-player game $N = \{1\}$ the Recursive Core is trivially defined.
2. *Recursion and dominance* Now assume that the Recursive Core has been defined for all games with $k - 1$ players. Then for a k -player game $N = Q \cup \overline{Q}$ an outcome $\omega = (x, P)$ is dominated through a coalition \overline{Q} forming partition $P^{\overline{Q}}$ and an outcome $(y, P^{\overline{Q}} \cup P^Q) \in \Omega(N, w)$, if
 - (a) when $\text{RC}(Q, w^{P^{\overline{Q}}}) \neq \emptyset$ then for all P^Q satisfying $(y^Q, P^Q) \in \text{RC}(Q, w^{P^{\overline{Q}}})$ and for all $i \in \overline{Q}$ it holds that $y^i > x^i$, and
 - (b) when $\text{RC}(Q, w^{P^{\overline{Q}}}) = \emptyset$ then for all P^Q and for all $i \in \overline{Q}$ it holds that $y^i > x^i$.
3. *Core* The Recursive Core of the PFF game (N, w) , denoted by $\text{RC}(N, w)$, is the set of undominated outcomes.

According to Definition 2.2 a deviating coalition \overline{Q} has to evaluate the profit it can achieve based on the outcome of the residual game, where a rational residual behavior is

expected. We assume conservatism of the deviators only among the remaining possible outcomes in the following sense. If the Recursive Core of the residual game is nonempty, then the members of coalition \bar{Q} expect the lowest possible payoffs of the *stable* outcomes $(y, P^{\bar{Q}} \cup P^Q) \in \Omega(N, w)$ with $(y^Q, P^Q) \in \text{RC}(Q, w^{P^{\bar{Q}}})$. If the Recursive Core of the residual game is empty, then the payoff of the members of the deviating coalition must be higher than the given one regardless of the coalition structure of the residual players.

The Recursive Core is well-defined, though it may be empty. Observe that all outcomes $\omega = (x, P) \in \text{RC}(N, w)$ are Pareto efficient; $\sum_{i \in (C \in P)} x^i = w(C, P)$ for all $C \in P$.

Based on the concept of the Recursive Core, a *minimal claim* function can be defined, which describes the minimal claim of each coalition in the corresponding PFF game reduced to that coalition. This function, termed v^{mc} in the following, may be applied in the same spirit as a characteristic function, since it assigns a unique value to each coalition, which they can secure for themselves if they deviated. The formal definition of v^{mc} is as follows.

Definition 2.3. Let us consider the residual game (\bar{C}, w^C) over the player set \bar{C} defined by the partition function $w^C(K, P^{\bar{C}}) = w(K, P^{\bar{C}} \cup C)$ where $K \in P^{\bar{C}} \in \mathcal{P}(\bar{C})$. Let us denote the Recursive Core of the residual game by $\text{RC}(\bar{C}, w^C)$. The minimal claim function v^{mc} can be defined as

$$v^{mc}(C) = \begin{cases} \min_{\sum_{i \in C} x^i} \{ \Omega(N, w) \mid (x, P^{\bar{C}}) \in \text{RC}(\bar{C}, w^C) \} & \text{if } \text{RC}(\bar{C}, w^C) \neq \emptyset \\ \min_{\sum_{i \in C} x^i} \{ \Omega(N, w) \} & \text{if } \text{RC}(\bar{C}, w^C) = \emptyset \end{cases}$$

where $v^{mc}(C)$ is the minimal claim of coalition C .

With the help of the minimal claim function, a characterization of the Recursive Core can be given as follows.

Lemma 2.4. *The Recursive Core $\text{RC}(N, w)$ of the game (N, w) is a collection of Pareto efficient outcomes $(x, P) \in \Omega(N, w)$, such that there is no coalition C with $v^{mc}(C) > \sum_{i \in C} x^i$.*

Proof. Holds by construction. □

Using the minimal claim function one can find the Recursive Core of the game very efficiently without needing to consider the deviations of all possible coalitions and partitions. We illustrate this through the 5-player game example of Kóczy (2006). We use the

simplified notation here following the original one, we write the coalitional values for a partition as a vector to keep the example as transparent as possible.

Example 2.5. Consider a game with $N = \{1, 2, 3, 4, 5\}$, and w given by;

$$w(123, 4, 5) = (10, 1, 1),$$

$$w(123, 45) = (0, 5),$$

$$w(12, 3, 4, 5) = (0, 1, 0, 0),$$

$$w(12, 3, 45) = (0, 1, 0),$$

$$w(1, 2, 3, 45) = (0, 0, 1, 0),$$

$$w(12, 345) = (5, 0),$$

$$w(1, 2, 345) = (1, 1, 8),$$

with all other payoffs being zero.

The minimal claim function can be derived as follows: The deviation of the coalition $C = \{1\}$ for instance induces the (only stable) partition $\tilde{P} = \{\{2\}, \{3, 4, 5\}\} \in \mathcal{P}(\{2, 3, 4, 5\})$ in the residual game $(\{2, 3, 4, 5\}, w^{\{1\}})$. Thus, the worth of Player 1 is $w(\{1\}, \tilde{P} \cup \{1\}) = 1$, hence $v^{mc}(1) = 1$. The minimal claims of the rest of the coalitions, shown in Table 1, can be derived in the same manner.

Table 1: Minimal claim function

C	$\{1\}$	$\{2\}$	$\{3\}$	$\{4\}$	$\{5\}$	$\{1,2\}$	$\{4,5\}$	$\{1,2,3\}$	$\{3,4,5\}$	other
$v^{mc}(C)$	1	1	0	1	1	0	0	0	0	0

Now, for a Recursive Core element it must be true that each coalition receives at least its worth as a payoff; $x^1, x^2, x^4, x^5 \geq 1$. This leaves us with two possible stable partitions; $P_1 = \{\{1, 2, 3\}, \{4\}, \{5\}\}$ with a total payoff of 12 and $P_2 = \{\{1\}, \{2\}, \{3, 4, 5\}\}$ with a total payoff of 10 to be divided appropriately.

3 Games with Uncertainty

We model uncertainty in a dynamic setup; we consider a game with two periods, $t \in T = \{0, 1\}$. In period 1, one state of nature s out of a finite set of states of nature S occurs. We define the state of nature for period 0 as state 0, so the set of all states is $S' = \{0\} \cup S$. Our

interest is clearly in the cases with non-degenerate uncertainty; $|S| > 1$. In period 1 the players are involved in a PFF-game $\Gamma_s = (N, w_s)$, where the game itself is allowed to be state-dependent. Period 0 serves as a point in time prior to the resolution of uncertainty.

Player $i \in N$ evaluates his payoffs by a utility function $u^i : \mathbb{R}^S \rightarrow \mathbb{R}$, which assigns to every profile of payoffs $x^i = (x_1^i, \dots, x_S^i) \in \mathbb{R}^S$ a utility level $u^i(x^i)$ and is assumed to be continuous and state-separable, i.e. $u^i(x^i) = \sum_{s \in S} u_s^i(x_s^i)$, where $u_s^i(x_s^i)$ is monotonically increasing. A well-known example of utility functions satisfying these assumptions is the von Neumann-Morgenstern utility function.

3.1 The Model

A PFF-game with uncertainty is defined as follows.

Definition 3.1. A partition function form game with uncertainty (PFU-game) Γ is a tuple (N, S', w, u) where $w = (w_1, \dots, w_S)$ and $u = (u^1, \dots, u^n)$.

Note that state 0 merely serves as a point in time where players face future uncertainty and may decide to cooperate (to pool risk for instance).

The *outcome* of the game is an ordered pair (x, P) , where the $x = (x^1, \dots, x^n) \in \mathbb{R}^{S \times N}$ matrix is called an *allocation*, and $P = (P_1, \dots, P_S)$ a partition for each subgame. Let $\Omega(x, P)$ denote the set of outcomes in Γ . The state- s component $x_s = (x_s^1, \dots, x_s^n) \in \mathbb{R}^N$ of an allocation is referred to as the allocation in state $s \in S$. The set of partitions in Γ_s is \mathcal{P}_s and the set of partitions in Γ_s of $C \in N$ is $\mathcal{P}_s(C)$ with a typical element P_s^C . Let Γ_s^R denote the residual PFF-game played in state s by player set R . The payoff allocation matrix must satisfy a feasibility constraint; given the partition P , for all $C \in P$ we have $\sum_{i \in C} x_s^i \leq w_s(C, P)$ for all $s \in S$. The central question of a PFU-game is finding a stable outcome.

We study which outcomes of the game Γ are stable. In general an outcome is stable if no coalition ever can benefit from a deviation. That is, we require that there should be no blocking possible in the subgames in $t = 1$ or before the resolution of the uncertainty in $t = 0$.

Thus we will require from a stable outcome (x, P) of the game Γ that each of its components (x_s, P_s) belongs to the Recursive Core of the subgame Γ_s to prevent any future deviation.

Now we are ready to define the Sustainable Core by means of the Recursive Core of suitably chosen subgames.

Definition 3.2 (Sustainable Core). An outcome (x, P) belongs to the Sustainable Core of the game Γ , denoted by $SC(\Gamma)$, if and only if

- (a) (x, P) is such that $(x_s, P_s) \in RC(\Gamma_s)$ for all $s \in S$,
- (b) and there is no coalition $R \in N$ forming some partition $\tilde{P}^R \in \mathcal{P}(\mathcal{R})$ and outcome $(\tilde{x}^R, \tilde{P}^R)$ such that $(\tilde{x}_s^R, \tilde{P}_s^R) \in RC(\Gamma_s^R)$ for all $s \in S$, and $u^i(\tilde{x}^i) > u^i(x^i)$ for all $i \in R$.

It means that for an outcome to belong to the Sustainable Core of the PFU-game Γ , the outcome should belong to the Recursive Core of the PFF-game Γ_s in every state $s \in S$. Moreover, no coalition (forming some partition) should be able to pick an element of the Recursive Core of the game restricted to R in every state, and in doing so improve utility in an ex ante sense. Since we require the deviating coalition to pick a Recursive Core element, no further counter-deviation from it can be expected by any sub-coalition; i.e. the blocking allocation is indeed self-enforcing.

Note, that if there is no externality, the game boils down to a transferable utility game with uncertainty, and so the definition of the Sustainable Core will coincide with that of the Weak Sequential Core. Similarly, if there is no uncertainty, than we have a single PFF-game with the Recursive Core as the solution to it.

In a PFU-game one can distinguish ex ante and ex post efficiency.

Definition 3.3. An outcome (\bar{x}, \bar{P}) is *ex ante efficient* in the game Γ if:

- (i) $\sum_{i \in N} \bar{x}^i \leq w(N, \bar{P})$, and
- (ii) there does not exist an outcome (x, P) with $\sum_{i \in N} x^i \leq w(N, P)$ such that $u^i(x^i) > u^i(\bar{x}^i)$ for all $i \in N$.

Definition 3.4. An outcome (\bar{x}, \bar{P}) is *ex post efficient* in the game Γ if $\sum_{i \in N} \bar{x}^i = w(N, \bar{P})$.

Note, that the concept of ex post efficiency says more than the usual feasibility conditions in TU-games, since it requires $\sum_{i \in N} \bar{x}_s^i = w_s(N, \bar{P})$ to hold at all states $s \in S$, but contrary to ex ante efficiency it does not imply Pareto efficiency, since it does not consider reallocation possibilities across states.

The next observation follows from the Pareto efficiency of the Recursive Core.

Corollary 3.5. *If $(\bar{x}, \bar{P}) \in \text{RC}(\Gamma)$ then (\bar{x}, \bar{P}) is ex post efficient.*

Note, on the other hand, that outcomes need not be ex ante efficient. Since binding agreements are not allowed for here, certain utility transfers across states might not be feasible.

4 Application: Safety Critical Rescheduling of Generators in The Electrical Power Transmission Network

The liberalization of the electricity markets, and the privatization and restructuring of power systems in the last decades resulted in an increased effort to fully economically exploit the infrastructure of the power grid. With this trend, the risk of failures has increased as demonstrated by the large blackouts observed (Fairley, 2004; Hines, J.Apt, and Talukdar, 2009). As emphasized by Beccuti, Demiray, Andersson, and Morari (2010), the situation is complicated by the fact that electrical power grids are large interconnected systems covering large areas where the control actions of a regional operator may adversely affect other components located hundreds of kilometers away. This observation suggests that the problem can be modeled by a PFF-game.

To avoid such dramatic consequences as country-wide blackouts, power system operators use various methodologies to keep the power grid in safe operational state. The protocols applied in these cases of security arrangements may affect the electricity market.

In this application we analyze, how the possibly occurring emergency procedures may be taken into account by the various stakeholders of the electricity market during the planning of transmission contracts. For the aim of simplicity we consider only line failures (no generator failures). We have to note that the method demonstrated here, the application of a PFU-game, can be straightforwardly applied to other scenarios of uncertainty; for example to network expansion scenarios, in which the uncertain expansion is assumed to be (partially) independent of the market participants, who represent the players of the transmission game.

4.1 PFF-game on The Transmission Network

Previously, a PFF-game for the electrical transmission networks has been defined in (Csercsik and Kóczy, 2011). The most important features of the applied transmission model are

the following. The admittance values (Y) and the injected/consumed energy amounts of the nodes uniquely determine the energy flows on the edges (branches or lines) of the network, which can be obtained by the solution of a system of linear equations. In addition to its admittance value, each edge is characterized by a transfer capacity (\bar{q}), which corresponds to the maximal amount of energy which can be transferred on it. Furthermore, as a most simple approach we neglect transmission losses. For the sake of simplicity we will assume that every node of the energy transmission network corresponds to a certain generator or consumer. For further information on DC load flow models, see Oren, Spiller, Varaiya, and Wu (1995) and Contreras (1997).

Players of the game correspond to nodes (generators or consumers), while coalitions of the game correspond to balancing groups for which the total inlet and outlet power has to be equal. A central independent network regulator may limit the power in and outlet of the coalitions (in other words reschedule the generators and curtail the loads) in order to maximize the total power transmitted by the network and secure the safe operation of the system and avoid line overloads. The value of a certain coalition in this setup is determined as twice the total transmission within the coalition.

In the current paper, based on the defined model, the safety-critical rescheduling of generators and load curtailment in the case of an instantaneous failure of a single transmission line is analyzed.

Electrical power transmission networks are safety critical systems, which have to be operated in a fault-tolerant manner. Assuming a basically safe operation, in the case of sudden line or generator failure, the flows in the network change instantaneously, potentially enhancing the vulnerability of the system. In such cases the rescheduling of the system is needed to prevent further, possibly more dramatic failures (Thanikachalam and Tudor, 1971; Kaltenbach and Hajdu, 1971).

As foreshadowed, in this work we will consider only line failures. Regarding our DC load flow model defined in (Csercsik and Kóczy, 2011) this means, that in the case of basic operation, the power in and outlets of the nodes have to be determined in a way, that if any line instantaneously fails, none of the remaining lines may be overloaded. We will call this a safe state of the system. This network stability requirement may be included in the LP problem described in eq. 13 of (Csercsik and Kóczy, 2011), by adding further inequality type constraints, which describe transfer limitations under line failures.

Basically, in the case of a line failure, the system may lose this stability property (the

failure of an additional more line may lead to network flows, which exceed the maximum capacity of a certain line, or lines). In this case the generators of the system need to be rescheduled and the loads are curtailed to bring the system in a safe state again. This must be done in a way which implies the minimal total change in generation and consumption values (Kaltenbach and Hajdu, 1971). In the following we will demonstrate the importance of the failure induced safety critical rescheduling on a simple 4-node network.

Example

Let us consider the network depicted in Figure 1.




Figure 1: The basic structure and parameters of the 4-node network. Y_{ij} corresponds to the admittance value of the line between i and j , while \bar{q}_{ij} denotes the value of the maximal possible energy transfer on the corresponding line (power transmission capacity). The numbers in parentheses at the nodes correspond to maximal generating capacity and desired consumption.

If we assume the grand coalition to form, which implies only one equality type constraint corresponding to the balance of total inlet and outlet power, the a safe network configuration corresponding to maximal total transmission (which in this case satisfies all consumers) is depicted in Figure 2a.

Since we assume a stable nominal operation, the line capacities are not exceeded in the case of the first failure (as depicted in Figure 2b), but without generator rescheduling and load curtailment after the second failure of line 2-4, line 2-3 is overloaded (which would lead to further failures). Via overheating of these lines, this may lead to further avalanche of failures and emergency shutdowns in critical case, which may lead to costs of billions of \$/EUR in most serious cases (Fairley, 2004). We assume that all generators can be




Figure 2: Basic (safe) operation of the 4 node network without failures, flows without rescheduling in the case of the failure of line 1-2, and in the case of the failure of line 1-2 and 2-4: The first failure does not lead to line overload, but without rescheduling the second failure causes the overload of line 2-3.

rescheduled, and all loads can be curtailed - of course this is not the case in a realistic scenario, where some consumers are treated with priority in general. If we reschedule the generators and apply load curtailment after the first failure of line 1-2 in order to bring the network in safe state again (we do this under the constraint of implying minimal total change in generation values and loads), the second failure of the line 2-4 will not cause overload of any other lines as depicted in Figure 3.




Figure 3: Rescheduling of the network after the failure of line 1-2: The second failure of line 2-4 does not overload the remaining lines. Line 2-3 operating on the edge of his capacity shows, that the rescheduling of generators and the curtailment of loads was justified.

4.2 The Power Transmission Game with Safety Critical Rescheduling

To define a cooperative game taking into account the uncertainty corresponding to the possible failure of lines, and the implied rescheduling which is causing changes in the transfer of certain coalitions, we use the formalism defined in Section 2.

In the case of m lines, each of which may fail, we have $S = m + 1$ number of states of nature in period $t = 1$. The first state, s_1 , corresponds to the normal (error-free) operation of the network, and the remaining m states describe the possible failures of the corresponding m lines. This implies that we will have $m + 1$ state-dependent partition function form games.

We will demonstrate the calculation of the Sustainable Core in the case of the network depicted in Figure 1. The state-dependent minimal claim function corresponding to the normal operation and to the failures of certain lines, and the implied safety critical rescheduling are listed in Table 1. $s = 1$ corresponds to the error-free operation of the network, while $s = 2$ corresponds to the error of the line 1-2, $s = 3$ to the error of the line 1-3, $s = 4$ to the error of the line 1-4, $s = 5$ to the error of the line 2-3, $s = 6$ to the error of the line 2-4 and $s = 7$ to the error of the line 3-4.

Table 2: Minimal claim function; v_s^{mc}

coalition	$s=1$	$s=2$	$s=3$	$s=4$	$s=5$	$s=6$	$s=7$
\emptyset	0	0	0	0	0	0	0
{1}	0	0	0	0	0	0	0
{2}	0	0	0	0	0	0	0
{3}	0	0	0	0	0	0	0
{4}	0	0	0	0	0	0	0
{1,2}	10	7	10	7	10	7	9.58
{1,3}	0	0	0	0	0	0	0
{1,4}	10	8	10	8	9.97	8	10
{2,3}	10	7	7	7	7.02	7	7
{2,4}	0	0	0	0	0	0	0
{3,4}	10	8	7	8	8	8	5.92
{1,2,3}	10	7	10	10	10	7	10
{1,2,4}	17.08	10	10	10	14.44	15	14.12
{1,3,4}	10	10	10	8	10	8	10
{2,3,4}	14.72	13	7	12	8	14.12	7
{1,2,3,4}	20	15	17	15	18	15	17

To evaluate the players' payoffs, we assume that all four players in our Example have

the following utility function;

$$u^i(x^i) = 0.88(1 - e^{-0.001x_1^i}) + \sum_{s=2}^{s=7} 0.02(1 - e^{-0.001x_s^i}) \text{ for all } i \in N.$$

One can interpret this expression as a von Neumann-Morgenstern utility function, where the probability of error-free operation is 88%, and the probability of any line failing is 2%.

Now consider the following allocation:

$$x = \begin{pmatrix} 2.91226 & 7.08774 & 2.91226 & 7.08774 \\ 2 & 5 & 2 & 6 \\ 5.7565 & 4.2435 & 2.7565 & 4.2435 \\ 3 & 4 & 3 & 5 \\ 5.51228 & 4.48772 & 3.21021 & 4.78979 \\ 0 & 7 & 0 & 8 \\ 5.53279 & 4.36166 & 2.63834 & 4.46721 \end{pmatrix} \in \mathbb{R}^{S \times N}.$$

It is easy to show that this allocation belongs to the Sustainable Core of the above defined game. First, one can check that $x_s \in \text{RC}(\Gamma_s)$ holds for all $s \in S$, thus there is no blocking possibility in time period $t = 1$. Now we only need to see if Part (b) of Definition 3.2 holds. We show next that x maximizes the sum of the players' utilities over allocations x^C with $x_s^C \in \text{RC}(\Gamma_s^C)$ for all $s \in S$.

Consider the following constrained maximization problem,

$$\begin{aligned} \max_{x^C} \sum_{i \in C} u^i(x^i) \\ \text{s.t. } \sum_{i \in C} x_s^i &= v_s^{mc}(C), \quad s \in S, & (1) \\ \sum_{i \in D} x_s^i &\geq v_s^{mc}(D), \quad s \in S, \emptyset \neq D \subsetneq C, & (2) \end{aligned}$$

where condition (1) is required for ex post efficiency and inequality (2) is a no-blocking condition. A solution to the maximization problem maximizes the sum of the players' utilities among those allocations that belong to $\text{RC}(\Gamma_s^C)$ for all $s \in S$. Since the given allocation x is a solution to this problem, it follows that there is no blocking possible in period $t = 0$ either.

Hence, $x \in \text{SC}(\Gamma)$.

It is worthwhile to note that this outcome is not really sensitive to the exact choice of the utility functions; e.g. the given x would also be a solution if the utility functions were

$$u^i(x^i) = 0.4(1 - e^{-0.01x_1^i}) + \sum_{s=2}^{s=7} 0.1(1 - e^{-0.01x_s^i}) \text{ for all } i \in N.$$

This example demonstrates that using a cooperative game and the concept of the Sustainable Core may help us to find such stable solutions to such difficult problems as the safety-critical operation of the power transmission networks that take into account possible line failures as well. This solution may be used in reality to facilitate immediate reaction to possible failures, which might lead to disastrous outcomes if reaching an agreement after the failure would take any time.

5 Conclusion

While in numerous real-life situations agreements have to be achieved in an environment with externalities and uncertainty about the future, the cooperative game theory literature has not considered this scenario yet. In this paper we introduced a new class of games, the partition function form games with uncertainty, to provide a framework to analyze these problems. We also proposed the Sustainable Core to solve the game, which is an extension of the Recursive Core and the Weak Sequential Core to this more complex setup.

As a possible application we analyzed the game of consumers and generators on a lossless DC load flow model of the electrical energy transmission network. While the described safety critical rescheduling of generators and curtailment of loads in the case of an arising line failure is a pure technological approach to avoid further overloads and shutdowns, dealing with its possibility in energy transmission contracts raises economical questions, for which the concept of the introduced sustainable core may serve as a possible solution concept. In this concept we assumed that the probabilities of line failures are known to each player of the game. A straightforward possible future scenario could be to analyze how asymmetric information affects the results. In most of the practical cases, regional generators have detailed information only about the network parameters in their area. Players who can obtain information about other areas may possibly effectively increase their bargaining potential and expected payoff compared to others.

The proposed framework may serve as a solution for many problems from environmental agreements to the formation of trading blocks or common pool resource allocation issues.

References

- AUMANN, R. J., AND B. PELEG (1960): “Von Neumann-Morgenstern Solutions to Cooperative Games Without Side Payments,” *Bulletin of the American Mathematical Society*, 66, 173–179.
- BECCUTI, A., T. DEMIRAY, G. ANDERSSON, AND M. MORARI (2010): “A Lagrangian Decomposition Algorithm for Optimal Emergency Voltage Control,” *Power Systems, IEEE Transactions on*, 25(4), 1769–1779.
- BESTER, H. (1984): “Core and equilibrium in incomplete markets,” *Journal of Economics*, 44(3), 255–266.
- CHANDER, P., AND H. TULKENS (1997): “The core of an economy with multilateral environmental externalities,” *International Journal of Game Theory*, 26(3), 379–401.
- CONTRERAS, J. (1997): “A Cooperative Game Theory Approach to Transmission Planning in Power Systems,” Ph.D. thesis, University of California, Berkeley.
- CSERCSEK, D., AND L. Á. KÓCZY (2011): “Externalities in the games over electrical power transmission networks,” IEHAS Discussion Papers 1125, Institute of Economics, Hungarian Academy of Sciences, Budapest.
- FAIRLEY, P. (2004): “The unruly power grid,” *IEEE Spectrum*, 41, 22–27.
- GILLIES, D. B. (1959): “Solutions to general non-zero-sum games,” in *Contributions to the Theory of Games IV*, ed. by A. W. Tucker, and R. D. Luce, no. 40 in *Annals of Mathematics Studies*, pp. 47–85. Princeton University Press, Princeton.
- GROSSMAN, S. J. (1977): “A characterization of the optimality of equilibrium in incomplete markets,” *Journal of Economic Theory*, 15(1), 1–15.
- HABIS, H., AND P. J. J. HERINGS (2010): “A Note on The Weak Sequential Core of Dynamic TU Games,” *International Game Theory Review*, 12, 407–416.
- (2011a): “Core Concepts for Incomplete Market Economies,” *Journal of Mathematical Economics*, 47(4-5), 595–609.

- HABIS, H., AND P. J.-J. HERINGS (2011b): “Transferable utility games with uncertainty,” *Journal of Economic Theory*, 146(5), 2126–2139.
- HART, S., AND M. KURZ (1984): “Stable Coalition Structures,” in *Coalitions and Collective Action*, ed. by M. Holler, pp. 236–258. Physica Verlag, Vienna.
- HINES, P., J. APT, AND S. TALUKDAR (2009): “Large blackouts in North America: Historical trends and policy implications,” *Energy Policy*, 37(12), 5249 – 5259.
- HUANG, C.-Y., AND T. SJÖSTRÖM (2003): “Consistent Solutions for Cooperative Games with Externalities,” *Games and Economic Behavior*, 43, 196–213.
- KALTENBACH, J., AND L. HAJDU (1971): “Optimal Corrective Rescheduling For Power System Security,” *IEEE Transactions on power Apparatus and Systems*, 90, 843–851.
- KÓCZY, L. Á. (2006): “A recursive core for partition function form games,” Research Memorandum RM/06/031, METEOR, Maastricht University, Maastricht.
- (2007): “A recursive core for partition function form games,” *Theory and Decision*, 63(1), 41–51.
- KOUTSOUGERAS, L. C. (1998): “A two-stage core with applications to asset market and differential information economies,” *Economic Theory*, 11(3), 563–584.
- KRANICH, L., A. PEREA, AND H. PETERS (2005): “Core concepts for dynamic TU games,” *International Game Theory Review*, 7(1), 43–61.
- MOULIN, H. (1985): “The Separability Axiom and Equal Sharing Methods,” *Journal of Economic Theory*, 36(1), 120–148.
- OREN, S., P. SPILLER, P. VARAIYA, AND F. WU (1995): “Folk Theorems on Transmission Access: Proofs and Counter Examples,” Working papers series of the Program on Workable Energy Regulation (POWER) PWP-023, University of California Energy Institute 2539 Channing Way Berkeley, California 94720-5180, www.ucei.berkeley.edu/ucei.
- PREDTETCHINSKI, A., P. J.-J. HERINGS, AND H. PETERS (2002): “The strong sequential core for two-period economies,” *Journal of Mathematical Economics*, 38, 465–482.

- REPULLO, R. (1988): “The Core of an Economy with Transaction Costs,” *Review of Economic Studies*, 55(3), 447–458.
- SHENOY, P. P. (1980): “A Dynamic Solution Concept for Abstract Games,” *Journal of Optimization Theory and Applications*, 32(2), 151–169.
- THANIKACHALAM, A., AND J. TUDOR (1971): “Optimal Rescheduling of Power for System Reliability,” *Power Apparatus and Systems, IEEE Transactions on*, PAS-90(5), 2186 – 2192.
- THRALL, R. M., AND W. F. LUCAS (1963): “ n -Person Games in Partition Function Form,” *Naval Research Logistics Quarterly*, 10(4), 281–298.
- VAN DER LAAN, G., AND M. CORNET (1998): “The Core of Bargaining Games with Externalities,” mimeo.